

## **The Canadian Entomologist**

**Index for Volume 138 (2006) / Index pour le volume 138 (2006)**



**Entomological Society of Canada /  
Société d'entomologie du Canada**

- Alfaro RI. See Kope *et al.*, 253  
Anderson GS. See Hobischak *et al.*, 376  
Anderson RS. See Zeran *et al.*, 123  
Ballantyne L. See Fu and Ballantyne, 339  
Barnfield P. See Gray *et al.*, 305  
Baum KA, Rubink WL, Coulson RN. Trapping of feral honey bee workers (Hymenoptera: Apidae) in a coastal prairie landscape: effects of season and vegetation type, 228  
Belton P. See Borkent and Belton, 91  
Bennett AMR. A new genus of Oedemopsini (Hymenoptera: Ichneumonidae: Tryphoninae) with a key to the genera of the tribe, 464  
Bennett RG. Ontogeny, variation, and synonymy in North American *Cybaeus* spiders (Araneae: Cybaeidae), 473  
Bergmann J. See Szöcs *et al.*, 263  
Berthiaume R. See Hébert *et al.*, 114  
Blossey B. See Dávalos and Blossey, 118  
Bordeleau C. See Hébert *et al.*, 114  
Borden JH. See Broberg *et al.*, 357  
Borkent A, Belton P. Attraction of female *Uranotaenia lowii* (Diptera: Culicidae) to frog calls in Costa Rica, 91  
Brauner AM, de Groot P. Field evaluation of host kairomones and pheromones for capture of *Conophthorus coniperda* (Coleoptera: Scolytidae), 723  
Broadbent AB. See Jandricic *et al.*, 712  
Broberg CL, Borden JH, Gries R. Olfactory and feeding preferences of *Cryptorhynchus lapathi* among hosts and nonhosts, 357  
Brown K. See Schlamp *et al.*, 384  
Cannings RA, Cannings SG. The Mantispidae (Insecta: Neuroptera) of Canada, with notes on morphology, ecology, and distribution, 531  
Cannings SG. See Cannings and Cannings, 531  
Cárcamo HA, Larson TR, Herle CE, Otani JK. Life history of *Lygus keltoni* (Hemiptera: Miridae) in the laboratory, 871  
Chan C-K. See Raworth *et al.*, 104  
Cline AR. See Majka and Cline, 314  
Cook J. See Majka *et al.*, 602  
Coulson RN. See Baum *et al.*, 228  
Currie DC. See Rowe *et al.*, 568  
Currie RW, Gatién P. Timing acaricide treatments to prevent *Varroa destructor* (Acari: Varroidae) from causing economic damage to honey bee colonies, 238  
Danks HV. Insect adaptations to cold and changing environments, 1  
Short life cycles in insects and mites, 407  
Dávalos A, Blossey B. Assessing capture techniques for ground beetles, 118  
de Groot P. See Brauner and de Groot, 723  
DeLury NC, Judd GJR, Gardiner MGT. Attraction of male *Pandemis limitata* (Lepidoptera: Tortricidae) to natural and synthetic pheromone sources: importance for assessing communication disruption, 697  
Despland E. See Noseworthy and Despland, 367  
Doddall L. See Gibson *et al.*, 285  
El-Sayed AM. See Trimble and El-Sayed, 218  
Fibiger M. See Lafontaine and Fibiger, 610  
Fields P. See Hou *et al.*, 95  
Floreni C, Pavan F, Nazzi F. Analysis of cuticular hydrocarbons in two *Anagrus* species (Hymenoptera: Mymaridae) as a tool to improve their correct identification, 348  
Footitt RG. See Scudder and Footitt, 24  
See Raworth *et al.*, 104  
Francke W. See Szöcs *et al.*, 263  
Fu X, Ballantyne L. *Luciola lei* sp. nov., a new species of aquatic firefly (Coleoptera: Lampyridae: Luciolinae) from mainland China, 339  
Fu X, Nobuyoshi O, Zhang Y, Lei C. A rearing apparatus and diet for the aquatic firefly *Luciola lei* (Coleoptera: Lampyridae), 399  
Fu X, Nobuyoshi O, Vencl FV, Lei C. Life cycle and behaviour of the aquatic firefly *Luciola lei* (Coleoptera: Lampyridae) from Mainland China, 860  
Galloway TD. See Mironov and Galloway, 165  
Gardiner MGT. See DeLury *et al.*, 697  
Garrison RW, von Ellenrieder N. Generic diagnoses within a closely related group of genera: *Brechmorhoga*, *Gynothemis*, *Macrothemis*, and *Scapania* (Odonata: Libellulidae), 269  
Gatién P. See Currie and Gatién, 238  
Germain C. See Klimaszewski *et al.*, 737

- Gibson GAP, Gillespie DR, Dosdall L. The species of Chalcidoidea (Hymenoptera) introduced to North America for biological control of the cabbage seedpod weevil, and the first recovery of *Stenomalina gracilis* (Chalcidoidea: Pteromalidae), 285
- Giffen D. See Olfert *et al.*, 875
- Gillespie DR. See Gibson *et al.*, 285
- Gillespie DR, Quiring DMJ. *Orius insidiosus* (Hemiptera: Anthocoridae) in the lower Fraser Valley, British Columbia, Canada, 590
- Grant GG. Foliage architecture explains oviposition preference of spruce budworm (Lepidoptera: Tortricidae) for white spruce over balsam fir, 198
- Gray DA, Barnfield P, Seifried M, Richards MH. Molecular divergence between *Gryllus rubens* and *Gryllus texensis*, sister species of field crickets (Orthoptera: Gryllidae), 305
- Gries G. See Schlamp *et al.*, 384
- Gries R. See Broberg *et al.*, 357
- See Schlamp *et al.*, 384
- Grixti JC, Packer L. Changes in the bee fauna (Hymenoptera: Apoidea) of an old field site in southern Ontario, revisited after 34 years, 147
- Hamilton KGA. The planthopper genus *Stenocranus* in Canada: implications for classification of Delphacidae (Hemiptera), 493
- Handfield D. See Handfield and Handfield, 333
- Handfield D, Handfield L. A new species of *Plusia* (Lepidoptera: Noctuidae) from North America, 853
- Handfield L. See Handfield and Handfield, 853
- Handfield L, Handfield D. A new species of *Capis* (Lepidoptera: Noctuidae) from Québec, Canada, 333
- Harris CR. See Jandricic *et al.*, 712
- Hart M. See Schlamp *et al.*, 384
- Hartley S. See Olfert *et al.*, 875
- Hébert C. See Klimaszewski *et al.*, 737
- Hébert C, Berthiaume R, Bordeleau C. Polyurethane foam strips to estimate parasitism of hemlock looper (Lepidoptera: Geometridae) eggs by *Telenomus* spp. (Hymenoptera: Scelionidae), 114
- Henderson D. See Kabaluk *et al.*, 647
- Henry TJ. Revision of the New World lygaeoid genus *Epipolops* (Heteroptera: Geocoridae: Pamphantinae: Epipolopini), with descriptions of five new species, 504
- Herle CE. See Cárcamo *et al.*, 871
- Hobischak NR, VanLaerhoven SL, Anderson GS. Successional patterns of diversity in insect fauna on carrion in sun and shade in the Boreal Forest Region of Canada, near Edmonton, Alberta, 376
- Hou X, Taylor W, Fields P. Effect of pea flour and pea flour extracts on *Sitophilus oryzae*, 95
- Huang M, Zhang Y. On the genus *Baaoira* (Hemiptera: Cicadellidae: Typhlocybinae) from China, with descriptions of five new species, 189
- Humble LM. Overwintering adaptations in Arctic sawflies (Hymenoptera: Tenthredinidae) and their parasitoids: cold tolerance, 59
- See Zilahi-Balogh *et al.*, 595
- Isman MB. See Wilson and Isman, 578
- Jackman JA. See Majka and Jackman, 292
- Jandricic S, Scott-Dupree CD, Broadbent AB, Harris CR, Murphy G. Compatibility of *Atheta coriaria* with other biological control agents and reduced-risk insecticides used in greenhouse floriculture integrated pest management programs for fungus gnats, 712
- Judd GGR. See Schlamp *et al.*, 384
- Judd GJR. See DeLury *et al.*, 697
- Kabaluk JT, Vernon RS, Henderson D. Population development of the green peach aphid and beneficial insects in potato fields in British Columbia, 647
- Klimaszewski J, Pelletier G, Germain C, Work T, Hébert C. Review of *Oxypoda* species in Canada and Alaska (Coleoptera, Staphylinidae, Aleocharinae): systematics, bionomics, and distribution, 737
- Kok LT. See Zilahi-Balogh *et al.*, 595
- Kope HH, Alfaro RI, Lavallée R. Virulence of the entomopathogenic fungus *Lecanicillium* (Deuteromycota: Hyphomycetes) to *Pissodes strobi* (Coleoptera: Curculionidae), 253
- Lafontaine JD, Fibiger M. Revised higher classification of the Noctuoidea (Lepidoptera), 610
- Lamb RJ. See Wise *et al.*, 638
- Langor DW. See Wesley *et al.*, 87
- Larson TR. See Cárcamo *et al.*, 871
- Lavallée R. See Kope *et al.*, 253
- Leggo JJ, Shorthouse JD. Development of stem galls induced by *Diplolepis triforma* (Hymenoptera: Cynipidae) on *Rosa acicularis* (Rosaceae), 661
- Modification of galls of *Diplolepis triforma* (Hymenoptera: Cynipidae) by the parasitoids *Eurytoma spongiosa* (Hymenoptera: Eurytomidae) and *Glyptomerus stigma* (Hymenoptera: Torymidae), 681
- Lei C. See Fu *et al.*, 860
- See Fu *et al.*, 399
- Majka CG, Cline AR. Nitidulidae and Kateretidae (Coleoptera: Cucujoidea) of the Maritime provinces of Canada. I. New records from Nova Scotia and Prince Edward Island, 314

- Majka CG, Jackman JA. The Mordellidae (Coleoptera) of the Maritime provinces of Canada, 292
- Majka CG, Selig G. *Lacconotus punctatus* and the family Mycteridae (Coleoptera) newly recorded in Atlantic Canada, 636
- Majka CG, Cook J, Westby S. Introduced Carabidae (Coleoptera) from Nova Scotia and Prince Edward Island: new records and ecological perspectives, 602
- Maw E. See Raworth *et al.*, 104
- McKenzie RIH. See Wise *et al.*, 638
- Mironov SV, Galloway TD. New and little-known species of feather mites (Acari: Analgoidea: Pteronyssidae) from birds in North America, 165
- Mittapalli O, Shukle RH, Wise IL. Identification of *mariner*-like elements in *Sitodiplosis mosellana* (Diptera: Cecidomyiidae), 138
- Mohan S, Sivakumar SS, Venkatesh SR, Raghavan GSV. Simple technique to increase the sensitivity of probe traps in detecting *Cryptolestes ferrugineus* (Coleoptera: Laemophloeidae) in stored wheat, 733
- Murphy G. See Jandricic *et al.*, 712
- Nazzi F. See Floreani *et al.*, 348
- Nobuyoshi O. See Fu *et al.*, 860
- See Fu *et al.*, 399
- Noseworthy MK, Despland E. How do primary nutrients affect the performance and preference of forest tent caterpillars on trembling aspen? 367
- Olffert O, Giffen D, Hartley S. Analysis of grasshopper (Orthoptera: Acrididae) population surveys in Saskatchewan: 1972–2004, 875
- Osono T. See Sugiura *et al.*, 390
- Otani JK. See Cárcamo *et al.*, 871
- Otvos IS. See Szöcs *et al.*, 263
- Packer L. See Grixti and Packer, 147
- Pavan F. See Floreani *et al.*, 348
- Pelletier G. See Klimaszewski *et al.*, 737
- Pollock DA. See Roughley *et al.*, 545
- Quiring DMJ. See Gillespie and Quiring, 590
- Raghavan GSV. See Mohan *et al.*, 733
- Raworth DA, Schade D. Life-history parameters and population dynamics of *Ericaphis fimbriata* (Hemiptera: Aphididae) on blueberry, *Vaccinium corymbosum*, 205
- Raworth DA, Chan C-K, Footitt RG, Maw E. Spatial and temporal distribution of winged aphids (Hemiptera: Aphididae) frequenting blueberry fields in southwestern British Columbia and implications for the spread of *Blueberry scorch virus*, 104
- Richards MH. See Gray *et al.*, 305
- Roland J. Effect of melanism of alpine *Colias nastes* butterflies (Lepidoptera: Pieridae) on activity and predation, 52
- Roughley RE, Pollock DA, Wade DJ. Biodiversity of ground beetles (Coleoptera: Carabidae) and spiders (Araneae) across a tallgrass prairie – aspen forest ecotone in southern Manitoba, 545
- Rowe L, Westlake KP, Currie DC. Functional significance of elaborate secondary sexual traits and their evolution in the water strider genus *Rheumatobates*, 568
- Rubink WL. See Baum *et al.*, 228
- Salom SM. See Zilahi-Balogh *et al.*, 595
- Schade D. See Raworth and Schade, 205
- Schiller AJ. See Szöcs *et al.*, 263
- Schlamp KK, Brown K, Gries R, Hart M, Gries G, Judd GGR. Diel periodicity of sexual communication in *Anarsia lineatella* (Lepidoptera: Gelechiidae), 384
- Scott-Dupree CD. See Jandricic *et al.*, 712
- Scudder GGE, Footitt RG. Alien true bugs (Hemiptera: Heteroptera) in Canada: composition and adaptations, 24
- Seifried M. See Gray *et al.*, 305
- Selig G. See Majka and Selig, 636
- Shorthouse JD. See Leggo and Shorthouse, 661
- See Leggo and Shorthouse, 681
- Shukle RH. See Mittapalli *et al.*, 138
- Sivakumar SS. See Mohan *et al.*, 733
- Spence JR. See Wesley *et al.*, 87
- Strong WB. Seasonal changes in seed reduction in lodgepole pine cones caused by feeding of *Leptoglossus occidentalis* (Hemiptera: Coreidae), 888
- Sugiura S, Yamazaki K. First record of a psilid fly feeding on cecidomyiid galls, 235
- Sugiura S, Yamazaki K, Osono T. Consequences of gall tissues as a food resource for a tortricid moth attacking cecidomyiid galls, 390
- Szöcs G, Otvos IS, Schiller AJ, Bergmann J, Francke W. *Cameraria gaultheriella* and *C. lobatiella* attracted in Canada to (E,Z)-8,10-tetradecadienal, the sex pheromone of the European *C. ohridella*, 263
- Taylor W. See Hou *et al.*, 95

- Trimble RM, El-Sayed AM. Effect of certain monounsaturated dodecene and tetradecene acetates and alcohols on electroantennogram response and pheromone-mediated trap catch of the obliquebanded leafroller, 218
- VanLaerhoven SL. See Hobischak *et al.*, 376
- VencI FV. See Fu *et al.*, 860
- Venkatesh SR. See Mohan *et al.*, 733
- Vernon RS. See Kabaluk *et al.*, 647
- von Ellenrieder N. See Garrison and von Ellenrieder, 269
- Wade DJ. See Roughley *et al.*, 545
- Wesley J, Williams DJM, Langor DW, Spence JR. Spruce beetle (Coleoptera: Scolytidae) parasitoids: cephalic morphology of larvae and a key to species (Hymenoptera: Braconidae, Pteromalidae), 87
- Westby S. See Majka *et al.*, 602
- Westlake KP. See Rowe *et al.*, 568
- Wheeler TA. See Zeran *et al.*, 123
- Whistlecraft JW. See Wise *et al.*, 638
- Williams DJM. See Wesley *et al.*, 87
- Wilson JA, Isman MB. Influence of essential oils on toxicity and pharmacokinetics of the plant toxin thymol in the larvae of *Trichoplusia ni*, 578
- Winchester NN. Ancient temperate rain forest research in British Columbia, 72
- Wise IL. See Mittapalli *et al.*, 138
- Wise IL, Lamb RJ, McKenzie RIH, Whistlecraft JW. Resistance to Hessian fly (Diptera: Cecidomyiidae) in a Canadian spring wheat cultivar, 638
- Work T. See Klimaszewski *et al.*, 737
- Yamazaki K. See Sugiura and Yamazaki, 235
- See Sugiura *et al.*, 390
- Zeran RM, Anderson RS, Wheeler TA. Sap beetles (Coleoptera: Nitidulidae) in managed and old-growth forests in southeastern Ontario, Canada, 123
- Zhang Y. See Huang and Zhang, 189
- See Fu *et al.*, 399
- Zilahi-Balogh GMG, Humble LM, Kok LT, Salom SM. Morphology of *Laricobius nigrinus* (Coleoptera: Derodontidae), a predator of the hemlock woolly adelgid, 595



